Problem B. Scapegoat

Input file: standard input
Output file: standard output

Aori is very careless so she is always making troubles. One day she does it again, with N big troubles! But this time she seems to be at ease because she has found M Inklings to take all the blames. Each trouble can be measured by a severity number a_i. Each trouble needs at least one Inkling to take the blame, and each Inkling can help Aori to take the blame for exactly one trouble. If two or more Inklings take the same trouble, they will take this blame together and discuss how to divide this trouble into.. some trivial things.. to reduce the pressure on each Inkling, as long as the total severity on Inklings is equal to the severity of this trouble.

Inklings are so warm so that Aori wants to think a way to let the variance of severity on each Inkling to be minimal. Could you help Aori make her scapegoats?

Formally, the variance of variables is the expectation of the squared deviation of a variable from its mean:

$$\sigma^2 = \frac{1}{n}\sum(x_i - \frac{1}{n}\sum x_j)^2$$

Input

The first line of the input gives the number of test cases, T. T test cases follow.

For each test case, the first line contains two integers N and M, where N is the number of troubles, and M is the number of Inklings. The next line contains N integers $a_1, a_2, ..., a_N$ representing the severity of the troubles that Aori makes.

Output

For each test case, output one line containing "`Case #x: y`", where `x` is the test case number (starting from 1) and `y` is the minimal variance.

`y` will be considered correct if it is within an absolute or relative error of 10^{-8} of the correct answer.

Limits

- $1 \le T \le 100$.
- $1 \le N \le M \le 2 \times 10^5$.
- $1 \le a_i \le 10000$.
- $\sum M \le 3 \times 10^6$.

Example

standard input	standard output
3 3 6 1 2 3 5 10 5 6 7 8 9 6 6 1 1 4 5 1 4	Case #1: 0.000000000000 Case #2: 0.400000000000 Case #3: 2.888888888889

Note

For the first sample, Aori can let one Inkling to take the first trouble's blame, two Inklings for the second, and three Inklings for the third. The severity on each Inkling is 1 unit, so their variance should be 0.