Problem C
Circles Are Far from Each Other

Time limit: 3 seconds

Problem Description

You are given n circles and two integer parameters k and ℓ. The i-th circle has radius r_i and $r_i \geq r_j$ holds for every $1 \leq i < j \leq n$. The task is to draw these n circles on a 2D plane so that the following conditions are simultaneously satisfied. Before proceeding to the conditions, let us recall that:

- Each circle is uniquely determined by its center O and radius r.
- A circle of radius r is defined as the set of all points whose distance from its center O is **exactly** r. All distances considered in this task are Euclidean.
- The interior region of a circle is defined as the set of all points whose distance from its center O is **less than** r.
- We say that one circle C *encloses* another circle C' if all points of C' lie inside the interior region of C.

You need to draw these n circles to satisfy all of the following conditions:

1. The centers of all circles are collinear.
2. The distance between any two centers is at most k.
3. No two circles intersect.
4. If a circle encloses two circles C and C', then either C encloses C' or C' encloses C.
5. The first ℓ circles **may or may not** be enclosed by other circles, whereas each of the remaining $n - \ell$ circles **must** be enclosed by at least one circle.

An arrangement that satisfies all the above conditions is called *feasible*. For a feasible arrangement $\mathcal{A}$, define its *quality* $d(\mathcal{A})$ as the minimum distance between any two points belonging to different circles.

If there exists at least one feasible arrangement for the given case, output the maximum possible quality among all feasible arrangements. If no feasible arrangements exist, output 0.




Figure C.1: A feasible arrangement for sample input 1.




Figure C.2: An arrangement that is **not feasible** for sample input 1.

Input Format

The first line contains three integers k, n, and ℓ, representing the maximum distance between centers, the number of circles to be drawn, and the number of circles that may or may not be enclosed by other circles, respectively.

The second line contains n integers $r_1, r_2, \ldots, r_n$, where r_i is the radius of the i-th circle.

Output Format

If no feasible arrangements exist, output a single integer 0.

Otherwise, output the maximum possible quality $d(\mathcal{A})$ among all feasible arrangements $\mathcal{A}$ in the following format:

- If $d(\mathcal{A})$ is an integer, output it as a single integer.
- If $d(\mathcal{A})$ is not an integer, output it in the rational form a/b such that $1 \le a, b \le 10^9$, $\gcd(a, b) = 1$, and $|d(\mathcal{A}) - a/b|$ is minimized. If multiple such forms satisfy the constraints, print any.

Technical Specification

- $1 \le k \le 10^9$
- $2 \le n \le 10^5$
- $1 \le \ell \le \min\{n, 200\}$
- $1 \le r_i \le 10^9$
- $r_i \ge r_j$ for every $1 \le i < j \le n$.

Sample Input 1

```
15 4 3
7 5 3 1
```

Sample Output 1

```
3
```

Explanation of Sample 1: A feasible arrangement is illustrated in figure C.1, where each circle and its center are shown in a unique color.

In this arrangement, the minimum distance between any two points belonging to different circles is 3, and the two points witnessing this distance are marked as red dots. Hence, the quality of this arrangement is 3, which is the maximum possible among all feasible arrangements for this case. Note that only circle 4 is required to be enclosed within another circle, while the remaining circles may or may not be enclosed.

Figure C.2 shows an arrangement that is **not feasible**. In this case, circle 1 encloses both circle 3 and circle 4, but neither of these two circles encloses the other, thereby violating condition 4.

Sample Input 2

```
14 6 1
7 5 4 3 2 1
```

Sample Output 2

```
1
```

Sample Input 3

```
14 2 2
5 4
```

Sample Output 3

```
5
```

Sample Input 4

```
22 4 4
4 4 1 1
```

Sample Output 4

```
10/3
```

Sample Input 5

```
13 3 3
6 3 1
```

Sample Output 5

```
4
```